

# FIRE EFFECTS ON SOIL AND VEGETATION IN A DEGRADED BRAZILIAN SAVANNA

COSTA, Yasmmin Tadeu <sup>a\*</sup> BARCELOS, Anna Carolina <sup>b</sup> RODRIGUES, Silvio Carlos <sup>c</sup>

(a) Phd Student in Geography. State University of Center West (UNICENTRO), Guarapuava (PR), Brazil. ORCID ID: <https://orcid.org/0000-0002-7050-7339>. CURRICULUM LATTES: <http://lattes.cnpq.br/4069992796148459>

(b) Msc in Geography. Federal University of Uberlândia (UFU), Uberlândia (MG), Brazil. ORCID ID: <https://orcid.org/0000-0001-5461-4508>. CURRICULUM LATTES: <http://lattes.cnpq.br/9360376324743490>

(c) Phd in Geography. Professor of the Federal University of Uberlândia (UFU), Uberlândia (MG), Brazil. ORCID ID: <https://orcid.org/0000-0002-5376-1773>. CURRICULUM LATTES: <http://lattes.cnpq.br/0563872406702918>

## (\*) CORRESPONDING AUTHOR

Address: Av. João Naves de Ávila, 2121. Bloco H Sala 1H16, CEP: 38400-000, Uberlândia (MG), Brazil. Phone: (+55)34-3291-6390  
E-mail: [silgel@ufu.br](mailto:silgel@ufu.br)

## ABSTRACT

This research consists of a comparative study through the controlled application of fire under different conditions of natural and exotic phytophysiognomy of the Cerrado vegetation area. The area of study is located in Cerrado recovery vegetation and degraded pasture areas. On this area, five experimental plots were established from the specific characteristic of its vegetation cover, slope and soil type (treatments) and connected to a Gerlach trough that accumulates the runoff and were weekly measured. Two artificial fires were applied to plots in an interval of 8 months in attempt to understand the effects of fire on runoff, sediment yield and regrowth of vegetation. The results show that runoff do not present differences between treatments through the period of study. Medium values of sediment yield for all plots vary from 6.70 g L<sup>-1</sup> to 57.24 g L<sup>-1</sup> showing that the plots which present high protection by vegetation create conditions to diminish the impact of precipitation on the surface.

**Keywords:** Soil Erosion; Experimental Plots; Fire.

## RESUMO / RÉSUMÉ

### EFEITOS DO FOGO NO SOLO E VEGETAÇÃO NA SAVANA BRASILEIRA DEGRADADA

Esta pesquisa consiste em estudo comparativo sobre a aplicação controlada de fogo em diferentes condições de vegetação exótica e natural de Cerrado. A área de estudo está localizada em áreas de Cerrado sobre uso de pastagem degradada e áreas em recuperação. O delineamento da pesquisa contou com cinco parcelas experimentais abrangendo características específicas de vegetação, declividade e tipo de solo, conectadas à uma calha coletora do escoamento resultante na área, e mensurados semanalmente. Duas queimadas artificiais foram aplicadas em um intervalo de 8 meses de modo a compreender o efeito do fogo sobre o escoamento superficial, perda de solo e rebrota da vegetação. Os resultados mostraram que não houve diferença de escoamento superficial entre os tratamentos durante o período de análise. Valores médios de produção de sedimento para todas as parcelas variaram entre 6.70 g L<sup>-1</sup> e 57.24 g L<sup>-1</sup>, mostrando que as parcelas que obtiveram intensa cobertura vegetal criaram condições para reduzir o efeito da precipitação sobre a superfície.

**Palabras claves:** Erosão do Solo; Parcelas Experimentais; Fogo.


### EFFECTOS DEL FUEGO SOBRE EL SUELO Y LA VEGETACIÓN EN UNA SAVANA BRASILEÑA DEGRADADA

Esta investigación consiste en un estudio comparativo sobre la aplicación de quemada controlada en distintas condiciones natural y exótica de la vegetación del Cerrado. El área de estudio se encuentra con vegetación del Cerrado en recuperación y extensiones en pasturas degradadas. En esta área, se han establecido cinco parcelas experimentales que abarcan diferentes tipologías vegetal, declividad y tipo de suelo, conectadas a una canaleta regredora de escurrimiento que se midió semanalmente. Se han aplicado en las parcelas dos quemadas artificiales en una gama de 8 meses con lo objetivo de entender los efectos del fuego en el escurrimiento, producción de sedimentos y reestablecimiento de la vegetación. Los resultados no presentaron diferencias en los escurrimientos superficiales entre los tratamientos durante el período de análisis. Los valores medios de la producción de sedimentos fueron de 6,70 g L<sup>-1</sup> 57,24 g L<sup>-1</sup> en todas las parcelas, mostrando así que en las áreas con mayor cobertura vegetal hay creado condiciones que minimizan el impacto de la lluvia sobre la superficie.

**Palabras clave:** Erosión del Suelo; Parcelas Experimentales; Fuego.

## Article history:

Received 28 May, 2018  
Accepted 20 June, 2018  
Publisher 15 July, 2018



## INTRODUCTION

The increase of soil erosion on Cerrado Biome is a real problem to the maintenance of surface equilibrium. Fire on Cerrado is a natural phenomenon, but the repetition of fire has been triggered by an anthropic action. Repeated fires intensify this problem with the removal of soil protection provided by vegetation, because it is supposed that after the repetition of fires in a small time interval, the regrowth of vegetation is not sufficient to protect the soil to erosion process. A higher amount of studies of fire effects on soil erosion in the Cerrado Biome was carried out after natural fires and a few experimental research use controlled fire to comprehend this process. Trying to understand it, a set of plots was constructed to study the ecosystem responses as changes in parameters linked to soil erosion after repeated fires.

Ryan (2002) describes fire as disturbance process, which is defined by its aspect of frequency, extent, seasonality, magnitude, and synergy that will affect the environment in different ways. The dynamic and severity of this process vary in time and space depending on the biophysical environment and location of the fire perimeter.

Ecosystem responses to fire depends of some functional processes that are altered by fire and include recolonization, regeneration and watershed hydrology processes, such as soil erosion or natural revegetation; Fire intensity, fire severity and burn severity are operationally manageable measures used for prediction of their dynamics after fires (KEELEY, 2009).

Soil physical properties, which are important in the soil erosion process, are affected by fire. According to Bento-Gonçalves and Vieira (2013), color, ash content, water repellency, changes in the structure, texture and porosity are the main factor of soil affected by fire.

Brazilian Cerrado species are much more tolerant to fire than forest species. Researches on natural fire on Cerrado shows low rates of mortality in tree and shrub, with rapid resprout of herbaceous layer after fire (ROBERTS, 2015; PIVELLO ET AL., 2010; DE ANDRADE; MIRANDA 2014). Environmental characteristics as mean annual precipitation, duration of dry season, soil fertility and disturbance are the key factors that determine the density of grasses, trees and shrubs in each Cerrado phytophysiology (HOFFMANN ET AL., 2003; HOFFMANN ET AL., 2012; RODRIGUES; BEZERRA, 2009). Multiple factors contribute to this difference in flammability, including microclimate, fuel amount and fuel type. Savannas are highly flammable and therefore typically subjected to frequent fire.

The previous revision of literature provides some dilemma about the real relationship between soil erosion effects and action of fires, due to the presentation by authors about results with relationship of soil erosion with fire occurrence and other studies with indications that ecological responses are not predictable from fire intensity and severity. We can add to this the almost inexistence of studies about this theme on Cerrado Biome, and understand that further research is needed to comprehend how to translate burn severity or fire severity measures and research into useful predictions about soil erosion.

From this information, the present study aims to understand and quantify the effects of fire on the increase or decrease of the erosive processes with reference to five characteristics of vegetation existing on degraded Cerrado area, based on the assumption that the burning and removal of vegetation that cover of topsoil favors the flow of soil sediment entrainment.

## STUDY AREA

The study area is located in the county of Uberlândia (Minas Gerais), Brazil at the Gloria Experimental Station located at the following UTM coordinates: 22S fuse, 794065 E and 7902595 N, with altitudes from 860 to 850 m above sea level (Figure 1). The soil at the experimental plots presents a sandy loam texture and derives from sandstone and siltstone from the Marília formation

(cretaceous). Some plots were built over degraded sites on exposures of subsoil material or BW soil. Table 1 presents the main information of the soil and other surface materials. The local climatic conditions are characterized by two well-defined seasons with rain concentration from November to March and a dry period from April to October both controlled by the Atlantic air mass or Continental air mass. Natural vegetation was replaced by pasture and agricultural practices, some areas were abandoned, and a regrowth of natural species creates a mosaic of land uses.




Figure 1- Location of study site and plots A, B, C, D and E. Being PA = Plot A, PB = Plot B, PC = Plot C, PD = Plot D, PE = Plot E.

Table 1- Physic-chemical characteristics of the studied plots subjected to different treatments of land.

Plots	Landscape Condition	Vegetation Characteristics	Soil Physical Properties			Soil Chemical Properties		Moisture Content	
			Sand	Silt	Clay	pH <sup>1</sup>	CEC <sup>2</sup>	Wet <sup>3</sup>	Dry <sup>3</sup>
				g Kg <sup>-1</sup>		Water	mmolc dm <sup>-3</sup>		%
A	Degraded Pasture	Grass	896	3	101	5,1	20,3	8,04	2,78
B	Degraded Pasture	Grass and Shrub	891	13	96	5	28,5	9,59	3,81
C	Flood Area	Grass	817	39	144	5,4	21,4	16,89	10,42
D	Degraded Area	Native Grass	836	3	161	5,1	20,2	8,51	3,47
E	Recovery Area	Grass and Tree	535	77	388	5,1	32,4	6,74	2,77

<sup>1</sup>Soil Reaction. <sup>2</sup>Cation Exchange Capacities. <sup>3</sup>Season (Wet – November to April; Dry – March to October).

## MATERIALS AND METHODS

### EXPERIMENTAL SETUP

For the assessments of fire effects on soil erosion, five plots of 1m<sup>2</sup> located in a degraded Cerrado environment were carried out, being differentiated by varying factors such as slope, soil drainage, type and density of vegetation and sediment yield after the action of controlled fire. In these plots

were also installed a Gerlach trough runoff sampler. In the plot base a gallon with 30 l of capacity collection of sediment and runoff were used. Each plot was delimited by galvanized sheets, with 10 cm deep into the ground and a height of 50 cm above the soil surface to avoid material from splash (Figure 2). Each treatment plot has differences on land cover, soil texture and chemical properties.

A superficial sample of soil was collected for a fertility index, particles density, bulk density and total porosity analysis. The main information about soil characteristics of each plot is presented in Table 2. The parameters analyzed were runoff, sediment yield, vegetation cover, soil moisture and precipitation. The data collection period was held from 12 March 2014 to 05 June 2015, and the application of controlled fire occurred on April 16 and November 11 of 2014. The two controlled fires in all the plots were considered of low severity because it is possible to observe the restoration and maintenance of a variety of ecological attributes that are generally viewed before the fires as stated by Neary et al. (2005).



Figure 2 - Plot structure.

Table 2 - Physico-chemical parameters for the study area.

Parameters		Unit of measurement	Methodology used	Frequency of collection	
Soil	Chemistry	pH	pH on Water –EMBRAPA		
		Moisture	%	Apparent dielectric constant of moist soil	
		Macronutrients	mmolc dm <sup>-3</sup>	EMBRAPA methods (1997)	Once
		CEC	mmolc dm <sup>-3</sup>	∑(K + Ca + Mg + (H+Al))	
	Physics	Texture	g kg <sup>-1</sup>	Pipette method	
	Runoff	Liters	Measurement at field plots		
Erosion	Soil Loss	g/L	Filtering and weight of samples	Weekly	
	Precipitation	mm	Pluviometric Station		
Vegetation	Surface Cover	%	Photo Comparasion		

## FIELD DATA COLLECTION AND LABORATORY ANALYSES

Field work was carried out once a week, with records of runoff, sediment yield collected at the Gerlach trough and at the gallons. Runoff was measured at field, and 1 l of homogenized water and sediment was collected for laboratory procedures of filtering and sediment weight. Superficial moisture determination at each plot was accomplished with the aid of a field moisture measurer. Determination of sediment weight eroded from the plots was obtained by filtering method. The weight of sediments retained in the filters were measured at a precision balance. Sediments collected at the Gerlach trough were also measured.

Soil samples were taken from a 0-5 cm depth to determine physico-chemical characteristics. The samples were stored in plastic bags until analysis. This material was used to perform laboratory analyses. The samples were air dried and screened to remove material ≥2mm. Soil texture was

measured using pipette method (EMBRAPA, 1997). Soil pH was measured on water. Chemical parameters of P (meq-1), K, Ca, Mg and Al were measured and fertility conditions were calculated.

Rainfall parameters were monitored by a station positioned at the same site of the experiments. The rainfall parameters recorded were rainfall intensity, total volume and total duration of each rainfall event. A set of vertical photography was performed each week after the data collection, which permits to identify and measure the land cover at each plot using the method adapted by Pinese Júnior et al. (2008).

## RESULTS

For the definition of the severity of burns was used the qualitative description of Keleey (2009), that takes into account the observation in the field and the resulting characteristics of fire in the environment.

Thus, the plot A in the first burning has some green leaves unburned, however is possible to see part of the organic matter consumed, classified as low severity fire. In the second firing, all the herbaceous layer was consumed in plot A, presenting a large amount of ash in the topsoil. It was possible to find small instances of white ash on the ground, but the overall characteristics of fire were not considered for classification. Accordingly, the second burning on plot A was moderate or severe burn (Figure 3).



Figure 3 - Effects of first (left) and of second burns (right) on plot A.

The first fire in plot B had not fully affected shrub species, with intact leaves and scorched part of the stem, but the grass species were hit and carbonized, being classified the burning as low severity. The second fire in the plot B also reached the vegetation more intensely, carbonizing nearly all the leaves of the shrub species and totally the herbaceous layer. These characteristics qualify as a fire of severe burning on moderate surface (Figure 4).



Figure 4 - Effects of first (left) and second burns (right) on plot B.

In plot C, the first burning carbonized most of vegetation cover, but still remained vegetation, which defines the fire as low severity. The second burn, in turn, eliminated all herbaceous, remaining on the soil an ash of the structure of dead grass. Thus, the second burning at plot C was also classified as moderate or severe over the surface. It is possible to realize the removal of a significant portion of the volume of ash comparing the two burning of vegetation. (Figure 5).



Figure 5 - Effects of first (left) and second burns (right) on plot C.

The first burning in the plot D was the same pattern as in plot A, reaching most of the herbaceous layer, but that there remains a considerable amount of green grasses, classifying the fire in this situation as a low severity. The second burning also occurred similarly seen in the plots already mentioned, however, small green leaves remained. This second fire was classified as moderate or severe burning over the soil surface. The small amount of ash present in the soil after firing was due to the slight quantity of biomass produced in the interval between fires during the dry period (S2) and also the effects of runoff and wind which removed previous ash from first fire. (Figure 6).



Figure 6 - Effects of first (left) and second burns (right) on plot D.

The fire in plot E also occurred with the same pattern of the four previous explanations. The first burning was classified as low severity by the vegetation permanence and the second as moderate or severe surface burning (Figure 7).



Figure 7 - Effects of first (left) and second burns (right) on plot E.

Rainfall pattern in the Cerrado Biome shows two distinct seasons, one from April to October and another from November to March. In both seasons, the medium temperatures are almost the same. The huge difference occurs in the rain distribution, with almost 90% of the rainfall concentrated in the summer, with peaks of precipitation in the period of December to March. Figure 8 demonstrates the pattern for the last 30 years and for 2014 total precipitation at Universidade Federal de Uberlândia Meteorological Station. In the year of 2014 some changes in this pattern have occurred, mostly for precipitation, which was 30% less than the historical medium pattern. For the rain station situated in the study area the values of 14 months were recorded for this study, with the amount being 18% less in comparison with the normal pattern of the area. This data show that a dry year occurred when compared to the local pattern, on the time interval of our study.




Figure 8 - Pattern of precipitation in the last 30 years at Uberlândia Station at UFU and recorded value in 2014.

For this study three stages were used to exam the results. The first stage (S1) was the pre-fire treatment that involved from February 2014 to 16 April 2014 that was marked as the end of the rainy season, in which the total record of precipitation was 231.4 mm. The second stage (S2) comprises the period after the first fire treatment and runs from 16 April 2014 to 11 November 2014, which was considered the dry season with 67.4 mm. The third stage (S3) started after the second fire treatment at 11 November with the new rain season, in which 871.1 mm of precipitation was recorded and comprises the entire 2014/15 rainy season. In the first stage 76 rainy events on 49 days were recorded with a peak of 63.11 mm and lowest value of 1.2 mm. In the S2 stage, 55 rainy events on 17 days were measured, with a peak of 50.10 mm, which represents 75% of the total rain in this stage, and the longest drought period of 6 weeks without precipitation. The S3 stage presented 275 rainy events distributed on 93 days and a high value of 176,5 mm and 0.3 mm to the lowest measure. In this stage some weeks presented no rain events. These data show a great difference between rain during the three stages, and points that the amount of rain is one of the key information to understand the evolution of land cover and the rates of regrowth of vegetation after fire, the runoff and the sediment yield behavior during the studied time.

Results about the percentage of land cover and regrowth of vegetation after fire treatments show two groups of behavior (Figure 9). The first group is compound by plots A, C and D with grass cover of the surface. The second group comprises plots B and E in which shrub and tree are the main vegetation. Before the first fire treatment the same trend was observed, in which the plot B registered highest mean value with 92.53% of cover while plot E shows the lowest value with 73.74% of surface cover. After the first fire treatment, no changes on the trend were observed because again the highest mean value was obtained in plot B with 68.71% and the lowest value was found on plot E with 34.82% of surface cover (Table 3). This values are compatible with the assumptions of Pivello et al. (2010) which indicate that fires on the beginning of the dry season and after individual rains are cooler than fires in the late dry season. These values obviously show the effects of




Figure 9 - Vegetal Cover Monthly mean value and Monthly Total Precipitation at the three stages of treatment.

fire and drought on the land cover percentage and regrowth of vegetation because they are inferior when we compared with the S1 and S3 in rainy seasons. In this way the trend was maintained after the second fire treatment when the highest and lowest values were obtained at the same plots as previously done, with highest value at plot B with 89.96% and the lower at plot E, with 20.54% of surface cover. Similar results were found by González-Pelayo et al. (2010) in studies carried out in a Mediterranean-shrub ecosystem.

Table 3 - Mean values of surface cover of three stages on the five plots treatments. N means the number of weeks.

Stage	S1 (n=7)					S2 (n=26)					S3 (n=19)				
Plot	A	B	C	D	E	A	B	C	D	E	A	B	C	D	E
Mean (%)	86.9	92.5	74.4	82.6	73.74	46.3	68.7	56.1	45.8	34.8	57.5	89.9	48.9	48.9	20,5

The highest mean values of runoff were measured at plot E in the third stage with 10.63 l and a peak of 14.45 l immediately after the second fire, while the lowest mean value was obtained at plot D, with 0.55 l. Plot E showed the highest value of runoff before the first fire treatment, with 2.03 l, while plot D showed the lowest values with 0.55 l. After the two fire treatments the responses were the same, where plot E continued with the highest values (Stage 2 = 4.90 l and Stage 3 = 10.63 l) and B with the lowest (Stage 2 = 0.96 l and Stage 3 = 3.06 l). Plot E presents an isolated tree with poor herbaceous strata, showing an almost sealed surface after the occurrence of the two fire treatments and the intense rains. Boix-Fayos et al. (2006) indicated that sealing of the surface together with stoniness and outcrop rock functioned as controlling mechanisms of soil erosion with high importance on runoff production. This situation was also noted by Campo et al. (2006) that observed increases of runoff and sediment transport by surface sealing. In addition, the coarse texture of surface material and the presence of gravels affects the runoff generation. This condition could explain the wash of the plot surface and contribute with the high index of runoff and the decrease of values of sediment yield.

Table 4 - Mean values of runoff of the three stages on the five plots treatments. n means the number of weeks.

Stage	S1 (n=8)					S2 (n=8)					S3 (n=20)				
Plot	A	B	C	D	E	A	B	C	D	E	A	B	C	D	E
Mean (%)	1.4	1.6	1.3	0.5	2.0	1.3	1.4	1.7	0.9	4.9	6.0	5.5	4.8	3.0	10.6

Runoff and sediment yield presents clearly three different stages during the research, all of them associated to the changes in the seasonal climatic conditions and synchronized to the fire treatments. Figure 10 shows the variations of runoff and sediment yield and it is possible to see lower values of both at the second stage. Likewise, it is also possible to see that in the second stage a synchronicity of responses happens between precipitation and runoff as shown in July 2014. Plot C, which is located in an area subject to seasonal flood, shows the lowest levels and low amplitudes of runoff in all the stages.




Figure 10 - Runoff and Soil Loss at three stages of treatment.

The result about sediment yield shows no noticeable differences occurring between plots A to E. When analyzed all periods of measurement, with 59 records, the highest value is on plot D with a mean value of 60.26 g/l, as the lowest is on plot B with a mean value of 5.08 g/l. For the measuring done before the first fire treatment, values ranging from 15.98 g/l on plot D while 5.08 g/l was found on plot E (Table 5). After the first fire all plots presented increases in mean values, with 19.58 g/l for plot D and 6.70 g/l for plot C, which indicates an increase on transport of sediment in dry conditions and low protection of the surface after rain showers. After the second fire an increment of the mean value occurs, as showed the highest value with 60.26 g/l in plot D and lowest value at plot C with 11.42 g/l. These results show that plot C located in seasonally flooded area show the lowest values of sediment yield and the highest values are achieved at plot D, where native grass is the main vegetation type and apparently is more affected by fire.

Table 5 shows the distribution of the sediment yield during the entire period of study. Visible differences between plots can be observed such as the constant lower values recorded at plot B (Shrub) while higher values found on plot D (native grass). This high protection to soil surface was observed in the study of Howes and Abrahams 2003 that indicate a great infiltration on low intensity rainfalls in a shrubland ecosystem. Also the maintenance of the shrub structure, which dies after the fire treatments but has a quickly resprouting and creates a group of new leafs as observed by (Fidelis et al. 2010) in field experiments with natural fires in grassland landscape with high shrub population in the south Brazil.

Table 5 - Mean values of sediment yield of 3 stages on the five plots treatments. n means the number of weeks.

Stage	S1 (n=7)					S2 (n=13)					S3 (n=23)				
	Plot	A	B	C	D	E	A	B	C	D	E	A	B	C	D
Mean (%)	10.5	8.4	10.4	15.9	5.0	9.9	7.9	6.7	19.5	13.2	57.2	8.8	11.4	60.2	49.7

## SOIL CHEMICAL PROPERTIES

The effects of two simulations of fire on the chemical properties of the soil were made with comparison of parameter based on their initial conditions, i.e. before Stage 1 (S1) and then after the second firing (S2) (Table 6). It was possible observe no changes in the evaluated parameters. Small variations were observed in the pH values and m (saturation aluminum). The pH values have as standard an increase in all the plots, while values of m spite of its high percentage changes, did not show a consistent pattern. These parameters could be affected by the occurrence of heavy rain which washing the surface without a real possibility of incorporate the elements in the ash that covered surfaces, as well as due to the low amount of organic matter in the second fire.

Table 6 - Changes of values of soil chemical properties. AF\* means after second fire.

Plot	Ph	P me <sup>-1</sup>	K	Ca	Mg	Al	CEC	V	m
	(water)	mg dm <sup>-3</sup>	mmolc dm <sup>-3</sup>					%	
A	5.1	0.9	15	3	0	6	25.4	21.20	44.90
A - AF*	5.3	2.7	11	4	0	4	20.3	13.40	61.80
B	5	0.9	19	3	0	4	23.5	12.30	58.80
B - AF*	5	1.1	18	3	0	5	28.5	14.90	53.30
C	5.1	1.1	15	5	1	4	21.4	20.60	57.70
C - AF*	5.4	1.1	16	3	1	6	21.4	29.90	38.50
D	4.9	1	11	3	1	6	26.3	20.80	45.50
D - AF*	5.1	0.8	9	3	1	4	20.2	16.30	56.10
E	5	0.9	12	3	1	7	30.3	13.60	47.60
E - AF*	5.1	1	15	3	1	4	32.4	14.20	60.20

## FINAL CONSIDERATIONS

Starting in a natural condition of surface protection five plots with 1m<sup>2</sup> were built in a degraded landscape in the Cerrado Biome, each one with different land cover. Two fires, one in the beginning of the dry season and another in the transition to the rainy season were used to understand the variations caused by this treatment. One consideration should be done about the precipitation values, which reach lower values in the research period in comparison to the local annual medium values.

The severity of the two fires applied in this study was similar. In the first simulation, all plots showed low severity, low consumption of organic matter, a fact that was attributed to the vegetative state of the plants, the high moisture content in the atmosphere and the soil humidity at the burn time, which was held at the end of the rainy season. The second burn was carried out at the end of the dry season, before the start of more concentrated rainfall. At this time the vegetation was poorly developed, with low soil and relative humidity.

The results about the pattern of land surface protection shows that shrub vegetation presents the highest protection to the surface during the entire study, and a plot with isolated tree and sparse grass cover shows the lowest values for surface cover. Plots with grass cover presents lower values than for shrub and higher than for isolated tree.

The evaluation on the runoff generation shows that seasonal conditions has a high influence on the responses of this parameter. Runoff presents an increase of values, with lower values in the first stage before the treatments and with higher values just after the second fire treatment, which means that the vegetation resprout does not reaches the same protection to the surface after repeated fires in a small interval of time.

Sediment yield does not show noticeable differences between plots and treatments during the study. The general trend was that of increase of values during the experimental stages, with higher values after the second treatment in the rainy season.

Despite these results, it is not possible to establish a direct relationship between the occurrence of the simulated fires and the variation of the runoff, because it is already naturally seasonally. Also changes on soil chemical parameters do not show visible changes or patterns of behavior on the results. Therefore, some further experiments must be done to try to evaluate the possibility of the occurrence of fire compared with the ecological responses like the case of the runoff and sediment yield, in view of the complexity of the phenomenon.

## ACKNOWLEDGE

The authors thank Fundação de Amparo a Pesquisa do Estado de Minas Gerais, project APQ-00231-16 and Conselho Nacional de Desenvolvimento Científico e Tecnológico for providing access to resources and financial support to accomplish the research (Project 302654/2015-1).

## BIBLIOGRAPHIC REFERENCE

DE ANDRADE, L.A.Z.; MIRANDA, H.S. The dynamics of the soil seed bank after a fire event in a woody savanna in central Brazil. *Plant Ecology*, 215(10), pp.1199–1209, 2014. Available at: <http://link.springer.com/10.1007/s11258-014-0378-z>.

BENTO-GONÇALVES, A.; VIEIRA, A. **Grandes incêndios florestais, erosão, degradação e medidas de recuperação dos solos**, Braga: Núcleo de Investigação em Geografia e Planeamento (NIGP), 2013.

BOIX-FAYOS, C. et al. Measuring soil erosion by field plots: Understanding the sources of variation. *Earth-Science Reviews*, 78(3–4), pp.267–285, 2006. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0012825206000766>.

- CAMPO, J. et al. Occurrence of soil erosion after repeated experimental fires in a Mediterranean environment. **Geomorphology**, 82(3–4), pp.376–387, 2006.
- EMBRAPA. **MANUAL DE MÉTODOS DE ANÁLISE DE SOLO. CENTRO NACIONAL DE PESQUISA DE SOLO. 2.ED.** RIO DE JANEIRO: EMBRAPA-CNPQ, 1997.
- FIDELIS, A. et al. Population biology and regeneration of forbs and shrubs after fire in Brazilian Campos grasslands. **Plant Ecology**, 211(1), pp.107–117, 2010.
- GONZÁLEZ-PELAYO, O. et al. Rainfall influence on plot-scale runoff and soil loss from repeated burning in a Mediterranean-shrub ecosystem, Valencia, Spain. **Geomorphology**, 118(3–4), pp.444–452, 2010. Available at: <http://dx.doi.org/10.1016/j.geomorph.2010.02.014>.
- HOFFMANN, W. a. et al. Fuels or microclimate? Understanding the drivers of fire feedbacks at savanna-forest boundaries. **Austral Ecology**, 37(6), pp.634–643, 2012.
- HOFFMANN, W. A.; ORTHEN, B.; DO NASCIMENTO, P. K.V. Comparative fire ecology of tropical savanna and forest trees. **Functional Ecology**, 17(6), pp.720–726, 2003. Available at: isi:000187184900002.
- HOWES, D. A; ABRAHAMS, A.D. Modeling runoff and runoff in a desert shrubland ecosystem, Jornada Basin, New Mexico. **Geomorphology**, 53(1–2), pp.45–73, 2003.
- KEELEY, J.E. Fire intensity, fire severity and burn severity: A brief review and suggested usage. **International Journal of Wildland Fire**, 18(1), pp.116–126, 2009.
- NEARY, D.G.; RYAN, K.C.; DE BANO, L.F. **Wildland Fire in Ecosystems: effects of fire on soils and water.** Fort Collins, CO: Gen. Tech. Rep., Rocky Mountain Research Station General Technical Report. Vol. 4. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, 2005.
- PINESE JÚNIOR, J.; CRUZ, L.; RODRIGUES, S. Monitoramento de erosão laminar em diferentes usos da terra, Uberlândia–MG. **Revista Sociedade & Natureza**, 20(2), pp.157–175, 2008. Available at: <http://www.seer.ufu.br/index.php/sociedadennatureza/article/viewFile/9532/5772> [Accessed August 18, 2014].
- PIVELLO, V.R. et al. Effect of fires on soil nutrient availability in an open savanna in Central Brazil. **Plant and Soil**, 337(1), pp.111–123, 2010.
- ROBERTS, S.J. **Tropical fire ecology**. 2(2001), pp.286–291, 2015.
- RODRIGUES, S.C.; BEZERRA, J.F.R. Study of matric potential and geotextiles applied to degraded soil recovery, Uberlândia (MG), Brazil. **Environmental Earth Sciences**, 60(6), pp.1281–1289, 2009. Available at: <http://link.springer.com/10.1007/s12665-009-0268-1> [Accessed August 20, 2014].
- RYAN, K.C. Dynamic interactions between forest structure and fire behavior in boreal ecosystems. **Silva Fennica**, 36(1), pp.13–39, 2002.